

Saclay-SPhT T02/110

PHASE STRUCTURE OF SUPERSYMMETRIC MODELS AT FINITE TEMPERATURE

Moshe Moshe^a and Jean Zinn-Justin^b

^a Department of Physics, Technion – Israel Institute of Technology,
Haifa, 32000 ISRAEL

^b CEA-Saclay, Service de Physique Théorique
F-91191 Gif-sur-Yvette Cedex, FRANCE

and

Institut de Mathématiques de Jussieu–Chevaleret, Université de Paris VII

ABSTRACT

We study $O(N)$ symmetric supersymmetric models in three dimensions at finite temperature. These models are known to have an interesting phase structures. In particular, in the limit $N \rightarrow \infty$ one finds spontaneous breaking of scale invariance with no explicit breaking. Supersymmetry is softly broken at finite temperature and the peculiar phase structure and properties seen at $T = 0$ are studied here at finite temperature.

(a) Supported in part by the Israel Science Foundation grant number 193-00 and the VPR fund for promotion of research.

(a) email: moshe@physics.technion.ac.il

(b) email: zinn@spht.saclay.cea.fr

1 Introduction

Bosons and fermions behave differently as they interact with an heat bath and supersymmetry is softly broken. Breakdown of supersymmetry at finite temperature has attracted much attention since the early interest in supersymmetry and involved much controversy on its consequences, appearance and phase structure [1,2]. Other related issues such as the restoration of broken internal symmetries at finite temperature supersymmetric theories were also debated [3]. More recently, there is a continued interest in the thermodynamics of supersymmetric Yang-Mills theory [4] and temperature effects on the minimal supersymmetric model [5].

Two supersymmetric models are studied in this paper at finite temperature, the $O(N)$ symmetric supersymmetric $\vec{\Phi}^4$ field theory, which is shown to have a peculiar phase structure in the large N limit, and a supersymmetric non-linear σ -model.

At zero temperature, it has been found in the past [6,7,8] that the large N limit of the scalar $\vec{\phi}^6$ theory in $d = 3$ dimensions and the supersymmetric $\vec{\Phi}^4$ theory [9] in $d = 3$ posses spontaneous breaking of scale invariance unaccompanied by explicit breaking. Here, we find out which of the peculiar properties of the phase structure seen at $T = 0$ in these models are maintained at finite temperature and how phase transitions occur as the temperature, rather than the coupling constant, is varied.

2 Supersymmetric models in the large N limit

A simple supersymmetric model, with one scalar superfield, in three euclidean dimensions at zero temperature is considered first. We then examine a supersymmetric non-linear σ -model.

2.1 The supersymmetric scalar field in three dimensions

We consider the $O(N)$ invariant action

$$\mathcal{S}(\Phi) = \int d^3x d^2\theta \left[\frac{1}{2} \bar{D}\Phi \cdot D\Phi + NU(\Phi^2/N) \right], \quad (2.1)$$

where Φ is an N -component vector: $\Phi(\theta) = \varphi + \bar{\theta}\psi + \frac{1}{2}\bar{\theta}\theta F$. Here, θ is a two-component Majorana spinor, φ is an N -component real scalar field, ψ is an N -component, two-component Majorana spinor and F is an N -component auxiliary field. $D = \partial/\partial\bar{\theta} - \not{\partial}\theta$ is the covariant derivative, the integration measure is $d^2\theta = \frac{i}{2}d\theta_2d\theta_1$ and $\bar{\theta}_\alpha\theta_\beta = \frac{1}{2}\delta_{\alpha\beta}\bar{\theta}\theta$. More details about the conventions used in the article can be found in the appendix.

In component form for a generic super-potential, one finds

$$\mathcal{S} = \int d^3x \left[-\frac{1}{2}\bar{\psi}\not{\partial}\psi + \frac{1}{2}(\partial_\mu\varphi)^2 - \frac{1}{2}U'(\varphi^2/N)\bar{\psi}\psi - U''(\varphi^2/N)(\bar{\psi}\varphi)(\varphi\psi)/N + \frac{1}{2}\varphi^2U'^2(\varphi^2/N) \right]. \quad (2.2)$$

The theory violates parity symmetry. Space reflection is equivalent to the change $U \mapsto -U$

One of the subjects which will be of interest here is spontaneous breaking of scale invariance. When the binding potential $U(\Phi^2)$ that binds bosonic and fermionic $O(N)$ quanta is tuned to a particular strength at which $O(N)$ singlet massless bound states are created, spontaneous breaking of scale invariance occurs and the resulting massless Goldstone particles appear as a supersymmetric multiplet of a dilaton and "dilantino". Though one may expect this as a result of non-renormalization of the coupling constant, in fact this happens also in the non-supersymmetric case. Explicit breaking of scale invariance will appear only at the next to leading order in $1/N$ as in the scalar case in the $d = 3$. At $N \rightarrow \infty$ flat directions in parameter space appear in the supersymmetric theory while maintaining the system's ground state energy at zero.

2.2 Superfield formulation of large N steepest descent method.

In the large N limit, a constraint on Φ^2 is introduced by adding to the initial action the term

$$\mathcal{S}_L = \int d^3x d^2\theta L(\theta) [\Phi^2(\theta) - NR(\theta)], \quad (2.3)$$

and integrating over the superfields $L(\theta) = M + \bar{\theta}\ell + \frac{1}{2}\bar{\theta}\theta\lambda$ and $R(\theta) = \rho + \bar{\theta}\sigma + \frac{1}{2}\bar{\theta}\theta s$. $N - 1$ components of Φ are first integrated out while keeping a test-component superfield $\Phi_1 \equiv \phi$. One finds

$$\mathcal{Z} = \int [d\phi][dR][dL] e^{-\mathcal{S}_N(\phi, R, L)} \quad (2.4)$$

with the large N action

$$\begin{aligned} \mathcal{S}_N = \int d^3x d^2\theta & \left[\frac{1}{2}\bar{D}\phi D\phi + NU(R) + L(\phi^2 - NR) \right] \\ & + \frac{1}{2}(N-1)\text{Str} \ln [-\bar{D}D + 2L]. \end{aligned} \quad (2.5)$$

The two saddle point equations, obtained by varying the superfields ϕ and R , are

$$2L\phi - \bar{D}D\phi = 0, \quad (2.6a)$$

$$L - U'(R) = 0. \quad (2.6b)$$

The third saddle point equation, obtained by varying L , involves the ϕ super-propagator given by the solution to the equation:

$$(-\bar{D}D + 2L(\theta)) \Delta(k, \theta, \theta') = \delta^2(\theta' - \theta),$$

that is,

$$\Delta(k, \theta, \theta') = \frac{[1 + \frac{1}{2}M(\bar{\theta}\theta + \bar{\theta}'\theta') - \frac{1}{4}(\lambda + k^2)\bar{\theta}\theta\bar{\theta}'\theta']}{k^2 + M^2 + \lambda} + \frac{\bar{\theta}[i\not{k} - M]\theta'}{k^2 + M^2}. \quad (2.7)$$

At coinciding θ arguments, one finds

$$\Delta(k, \theta, \theta) = \frac{1 + M\bar{\theta}\theta}{k^2 + M^2 + \lambda} - \frac{M\bar{\theta}\theta}{k^2 + M^2}. \quad (2.8)$$

The trace over the group indices yields a factor $(N - 1)$. The third saddle point equation, obtained by varying L in Eq. (2.5), is thus (for $N \gg 1$)

$$\begin{aligned} R - \phi^2/N &= \frac{1}{N} \text{tr} \Delta(k, \theta, \theta) \\ &= \frac{1}{(2\pi)^3} \int d^3k \left[\frac{1 + M\bar{\theta}\theta}{k^2 + M^2 + \lambda} - \frac{M\bar{\theta}\theta}{k^2 + M^2} \right]. \end{aligned} \quad (2.9)$$

It is now convenient to introduce a notation for the boson mass:

$$m_\varphi \equiv m = \sqrt{M^2 + \lambda}. \quad (2.10)$$

and write the saddle point equations in component form. Eq. (2.6a) implies

$$F - M\varphi = 0, \quad (2.11a)$$

$$\lambda\varphi + MF = 0. \quad (2.11b)$$

Eliminating F between the two equations one finds $\varphi m^2 = 0$ and, thus, if the $O(N)$ symmetry is broken the boson mass m_φ vanishes. Eq. (2.6b) yields

$$M = U'(\rho), \quad (2.12a)$$

$$\lambda = m^2 - M^2 = sU''(\rho). \quad (2.12b)$$

Eq. (2.9) in component form is

$$\rho - \varphi^2/N = \frac{1}{(2\pi)^3} \int \frac{d^3p}{p^2 + m^2} \quad (2.13a)$$

$$s - 2F\varphi/N = \frac{2M}{(2\pi)^3} \int d^3p \left(\frac{1}{p^2 + m^2} - \frac{1}{p^2 + M^2} \right). \quad (2.13b)$$

In these equations, a cut-off Λ is being introduced explicitly. We set

$$\frac{1}{(2\pi)^d} \int^\Lambda \frac{d^d k}{k^2 + m^2} \equiv \Omega_d(m). \quad (2.14)$$

Below, we need the first terms of the expansion of $\Omega_d(m)$ for $m^2 \rightarrow 0$. Introducing the cut-off dependent constant

$$\rho_c = \frac{1}{(2\pi)^3} \int^\Lambda \frac{d^3 p}{p^2} = \Omega_3(0), \quad (2.15)$$

we rewrite Eqs. (2.13) as

$$\rho - \varphi^2/N = \rho_c - \frac{1}{4\pi} m, \quad (2.16a)$$

$$s - 2F\varphi/N = \frac{1}{2\pi} M (|M| - m). \quad (2.16b)$$

Note that, here, the change $U \mapsto -U$ corresponds to

$$F \mapsto -F, \quad s \mapsto -s, \quad M \mapsto -M.$$

Finally, we calculate the action density $\mathcal{E}$ corresponding to the action $\mathcal{S}_N$ (Eq. (2.5)), $\mathcal{E} = \mathcal{S}_N/\text{volume}$, for vanishing fermion fields:

$$\begin{aligned} \mathcal{E}/N = & -\frac{1}{2} F^2/N + \frac{1}{2} s U'(\rho) + M F \varphi/N + \frac{1}{2} \lambda \varphi^2/N - \frac{1}{2} M s - \frac{1}{2} \lambda \rho \\ & + \frac{1}{2} \text{tr} \ln(-\partial^2 + M^2 + \lambda) - \frac{1}{2} \text{tr} \ln(\not{\partial} + M). \end{aligned} \quad (2.17)$$

In $d = 3$, we have in Eq. (2.17):

$$\frac{1}{2} \text{tr} \ln(-\partial^2 + M^2 + \lambda) - \frac{1}{2} \text{tr} \ln(\not{\partial} + M) = \frac{1}{2} \rho_c \lambda - \frac{1}{12\pi} (m^3 - |M|^3). \quad (2.18)$$

Note that the saddle point equations are recovered, in component form, by taking the derivatives of $\mathcal{E}$ with respect to the various parameters.

Using the saddle point Eqs. (2.6a), (2.16) and (2.12a) to eliminate F, s, ρ , one eliminates also the explicit dependence on the super-potential U and the expression simplifies into

$$\mathcal{E}/N = \frac{1}{2} M^2 \varphi^2/N + \frac{1}{24\pi} (m - |M|)^2 (m + 2|M|). \quad (2.19)$$

In this form we see that the action density $\mathcal{E}$ is positive for all saddle points, and, as a function of m , has an absolute minimum at $m = |M|$, and thus $\lambda = 0$, that is, for a supersymmetric ground state.

Eq. (2.6) implies $M\varphi = 0$, and thus $\mathcal{E} = 0$. Therefore, if a supersymmetric solution exists, it will have the lowest possible ground state energy and any non-supersymmetric solution will have a higher energy. Since M , m , and φ are related by the saddle point equations, it remains to verify whether such a solution indeed exists.

In the supersymmetric situation the saddle point equations reduce to $s = 0$, $M\varphi = 0$ and

$$\rho - \rho_c = \varphi^2/N - |M|/4\pi, \quad M = U'(\rho).$$

In the $O(N)$ symmetric phase $\varphi = 0$ and $|M| = 4\pi(\rho_c - \rho) = |U'(\rho)|$. In the broken phase $U'(\rho) = 0$ and $\varphi^2/N = \rho - \rho_c$. We will show in the next section that these conditions can be realized by a quadratic function $U(\rho)$ and, then, in both phases the ground state is supersymmetric and $\mathcal{E}$ vanishes.

2.3 The Φ^4 super-potential in $d = 3$: phase structure

We now consider the special case

$$U(R) = \mu R + \frac{1}{2}uR^2 \Rightarrow U'(R) = \mu + uR. \quad (2.20)$$

The dimensions of the θ variables and the field Φ are $[\theta] = -\frac{1}{2}$, $[\Phi] = \frac{1}{2} \Rightarrow [u] = 0$. Power counting thus tells us that the model is renormalizable in three dimensions. Prior to a more refined analysis, one expects coupling constant and field renormalizations (with logarithmic divergences) and a mass renormalization with linear divergences. Using the solution for the two-point function $W^{(2)}$, in Eq. (A1.6) in the appendix, one infers that the coefficient $A(p^2)$ has at most a logarithmic divergence, which corresponds to the field renormalization, while the coefficient $C(p^2)$ can have a linear divergence which corresponds to the mass renormalization.

For the quartic potential, Eqs. (2.12) are now

$$M = \mu + u\rho, \quad \lambda = us. \quad (2.21)$$

We introduce the critical value of μ :

$$\mu_c = -u\rho_c. \quad (2.22)$$

Taking into account Eqs. (2.21), one finds that Eqs. (2.16) can now be written as

$$M = \mu - \mu_c + u\varphi^2/N - \frac{u}{4\pi}\sqrt{M^2 + \lambda}, \quad (2.23a)$$

$$\lambda = 2uM\varphi^2/N + \frac{u}{2\pi}M \left(|M| - \sqrt{M^2 + \lambda} \right). \quad (2.23b)$$

Eqs. (2.23) relate the fermion mass $m_\psi = |M|$, the boson mass $m_\varphi = \sqrt{M^2 + \lambda}$ and the classical field φ . The phase structure of the model is then described

by the lowest energy solutions of these equations in the $\{\mu - \mu_c, u\}$ plane. The invariance under the change $U \mapsto -U$ that was mentioned above and seen now in Eq. (2.20) and the equations that followed. This invariance is reflected into the phase structure of the model and one can restrict the discussion to $u > 0$. The phase structure is summarized in Fig. 1 in the first and second quadrant of the $\{\mu - \mu_c, u/u_c\}$.

We find, indeed, that supersymmetry is left unbroken ($\lambda = 0$) and the ground state energy $\mathcal{E} = 0$ in each quadrant in the $\{\mu - \mu_c, u\}$ plane. This is consistent with Eqs. (2.23) having a common solution with $\lambda = 0$ (thus $m_\psi = m_\varphi = |M|$) and $M\varphi = 0$. They then reduce to

$$M = \mu - \mu_c + u \frac{\varphi^2}{N} - \frac{u}{4\pi} |M|, \quad M\varphi = 0. \quad (2.24)$$

The broken $O(N)$ symmetry phase: The $M = 0$ solution implies a spontaneously broken $O(N)$ symmetry, scalar and fermion $O(N)$ quanta are massless and

$$\varphi^2 = -N(\mu - \mu_c)/u, \quad (2.25)$$

which implies that this solution exists only for $\mu < \mu_c$. The solution exists in the fourth (and second) quadrant of the $\{\mu - \mu_c, u\}$ plane.

The $O(N)$ symmetric phase: Here, $\varphi^2 = 0$ and the equation

$$M = \mu - \mu_c - (u/u_c) |M| \quad (2.26)$$

yields the common mass M for the fermions and bosons. In Eq. (2.26) we have introduced the special value u_c of the coupling u :

$$u_c = 4\pi. \quad (2.27)$$

Eq. (2.26) splits into two equations, depending on the sign of M . The first solution

$$M = M_+ = (\mu - \mu_c)/(1 + u/u_c) > 0,$$

exists only for $\mu > \mu_c$ (first quadrant), as one would normally expect.

The second solution

$$M = M_- = (\mu - \mu_c)/(1 - u/u_c) < 0,$$

is very peculiar. There are two different situations depending on the position of u with respect to u_c :

(i) $u > u_c = 4\pi$ and then $\mu > \mu_c$: the solution is degenerate with another $O(N)$ symmetric solution M_+ .




Fig. 1 Summary of the phases of the model in the $\{\mu - \mu_c, u\}$ plane. Here $m_\varphi = m_\psi = |M_\pm| = (\mu - \mu_c)/(u/u_c \pm 1)$. The lines $u = u_c$ and $\mu - \mu_c = 0$ are lines of first and second order phase transitions.

(ii) $u < u_c$ and then $\mu < \mu_c$: the solution is degenerate with a solution of broken $O(N)$ symmetry.

The following different phases appear in the phase structure that is summarized in Fig. 1:

Region I : $\mu - \mu_c \geq 0$, $u/u_c \leq 1$:

Here, there is only one $O(N)$ -symmetric, supersymmetric ground state with $m_\psi = m_\varphi = M_+ = (\mu - \mu_c)/(u/u_c + 1)$ and $\varphi^2 = 0$.

Region II : $\mu - \mu_c \geq 0$, $u/u_c \geq 1$:

There are two degenerate $O(N)$ -symmetric ($\varphi^2 = 0$), supersymmetric ground states with masses $m_\psi = m_\varphi = M_+ = (\mu - \mu_c)/(u/u_c + 1)$ and $m_\psi = m_\varphi = -M_- = (\mu - \mu_c)/(u/u_c - 1)$.

Region III : $\mu - \mu_c \leq 0$, $u/u_c \geq 1$:

There is one supersymmetric ground state, it is an ordered state with broken $O(N)$ symmetry ($\varphi^2 \neq 0$, $m_\psi = m_\varphi = 0$).

Region IV : $\mu - \mu_c \leq 0$, $u/u_c \leq 1$:

There are two degenerate ground states: An $O(N)$ -symmetric , supersymmetric ground states with masses $m_\psi = m_\varphi = M_- = (\mu - \mu_c)/(u/u_c - 1)$ and $\varphi^2 = 0$. The second ground state is a supersymmetric, broken $O(N)$ symmetry state with $m_\psi = m_\varphi = 0$ and $\varphi^2 \neq 0$.

The action density: In order to exhibit the phase structure in terms of the variation of the action density $\mathcal{E}$, we plot in Fig. 2 the expression (2.19), but

use only the fermion gap equation in Eq. (2.23a), in such way that $\mathcal{E}$ remains a function of φ and λ , or equivalently φ and $m = \sqrt{M^2 + \lambda}$:

$$\frac{1}{N}\mathcal{E}(m, \varphi) = \frac{1}{2}M^2(m, \varphi)\frac{\varphi^2}{N} + \frac{1}{24\pi} [m - |M(m, \varphi)|]^2 (m_\varphi + 2|M(m, \varphi)|). \quad (2.28)$$




Fig. 2 **Region IV** : The energy density $W(m, \varphi) \equiv \frac{1}{N}\mathcal{E}(m, \varphi)$ as given in Eq. (2.28) as a function of the boson mass (m) and A , where $A^2 = \varphi^2/u_c$. Here $\mu - \mu_c = -1$, $u/u_c = 0.2$. As seen here there are two distinct degenerate phases. One is an ordered phase ($\varphi \neq 0$) with a massless boson and fermion, the other is a symmetric phase ($\varphi = 0$) with a massive ($m = |M_-|$) boson and fermion.

Fig. 3 displays the action density $\mathcal{E}$ restricted to $\varphi = 0$:

$$\frac{1}{N}\mathcal{E}(m, \varphi = 0) = \frac{1}{24\pi} [m - |\mu - \mu_c - (u/u_c)m|]^2 (m + 2|\mu - \mu_c - (u/u_c)m|). \quad (2.29)$$

Several peculiar phase transitions can be easily traced now in Eq. (2.28). First, one notes the phase transitions that occur when $\mu - \mu_c$ changes sign. When $0 < u < u_c$ and $\mu > \mu_c$, the system has a non-degenerate $O(N)$ -symmetric ground state with bosons and fermions of mass $M = M_+$. As $\mu - \mu_c$ changes sign ($0 < u < u_c$ fixed), two degenerate ground states appear, as seen in Fig. 2. Either $M = 0$ and $\varphi^2 = -N(\mu - \mu_c)/u$ or the system stays in an $O(N)$ -symmetric ground state with a mass $|M_-|$ for the bosons and fermions. Similarly, when one goes from $\mu < \mu_c$ to $\mu > \mu_c$ at $u > u_c = 4\pi$, the $O(N)$ symmetry is restored but there are two degenerate ground states to choose from, $M = M_\pm$.




Fig. 3 The action density $W(m) \equiv \frac{1}{N}\mathcal{E}(m, \varphi = 0)$ from Eq. (2.29) in region **II** of Fig. 1. Here $\mu - \mu_c = 1$ (sets the mass scale) and u/u_c varies between 1.6 and 1.2. There are two massive, degenerate $O(N)$ symmetric SUSY vacua with $\varphi = 0$ and possibly $|M_-/M_+| \gg 1$ (as $u \rightarrow u_c$) with $m_\psi = m_\varphi = M_+ = (\mu - \mu_c)/(u/u_c + 1)$ and $m_\psi = m_\varphi = -M_- = (\mu - \mu_c)/(u/u_c - 1)$.




Fig. 4 As in Fig.3, the action density $W(m) \equiv \frac{1}{N}\mathcal{E}(m, \varphi = 0)$ with $\mu - \mu_c = 1$ and $u/u_c = 1.2$ is changed to $u/u_c = 0.8$ (from region **II** to region **I** of Fig. 1). There are two degenerate $O(N)$ symmetric SUSY vacua at $u/u_c = 1.2$ with masses $m_\psi = m_\varphi = |M_\pm|$ where $|M_\pm| = \mu/(u/u_c \pm 1)$ while at $u/u_c = 0.8$ there is a non-degenerate vacuum at $m_\psi = m_\varphi = M_+$.

As seen in Figs. 3 and 4, an unusual transition takes place when one varies the coupling constant u . For positive $\mu - \mu_c$ we find two degenerate ground states if $u > u_c$. As u is lowered (at fixed $\mu - \mu_c$) the ground state with mass $m_\varphi = m_\psi = -M_-$ disappears ($|M_-/M_+| \rightarrow \infty$) and only the $O(N)$ -symmetric phase remains with $M = M_+$. Namely, suppose we consider at $\{u > u_c, \mu - \mu_c > 0\}$ a physical system in a state denoted by A and defined by $\{\varphi^2 = 0, M = M_-\}$. Such a system will go into a state B defined by $\{\varphi^2 = 0, M = M_+\}$ when u **decreases** and passes the value $u = u_c = 4\pi$. Now, if we consider the reversed process;

a physical system at $\{u < u_c, \mu - \mu_c > 0\}$ that is initially in the ground state B and u **increases** and passes $u = u_c$. There is now no reason for the system to go through the reversed transition from B to A since the $O(N)$ -symmetric states with $M = M_-$ and $M = M_+$ are degenerate and since supersymmetry is preserved the energy of state A will not decrease below zero. These peculiar phase transitions with $(|M_-/M_+| \rightarrow \infty)$ and with “infinite hysteresis” in the $A \rightarrow B$ transitions are due to the fact that supersymmetry is left unbroken and the degeneracy of the two possible ground states in the leading order in $1/N$. If supersymmetry would have been broken by some small parameter, the lifted degeneracy would, most probably, have translated into a slow first order transitions between the otherwise degenerate ground states.

The transition from the degenerate vacua at $u/u_c = 1.2$ to a non-degenerate ground state at $u/u_c = 0.8$ is shown in Fig. 4 (from region **II** to region **I**). This type of transitions between the different phases of Fig. 1 will be studied as a function of the temperature in the next section.

Special situation. In general, when $\mu = \mu_c$ the mass M vanishes. However, there is a special case when

$$u = u_c = 4\pi.$$

Then the value of M is left undetermined. An accumulation point of coexisting degenerate ground states exist in the phase structure shown in Fig. 1. The case $\mu = \mu_c$ represents a scale invariant theory where, however, the $O(N)$ fermionic and bosonic quanta can have a non-vanishing mass $m_\varphi = m_\psi = |M|$. Since u has not gone any perturbative renormalization there is no explicit breaking of scale invariance at this point. Thus, the only scale invariance breaking comes from the solution of the gap equation for M which leaves, however, its numerical value undetermined. We see a dimensional transmutation from the dimensionless coupling u that is fixed at a value of $u = u_c$ into an undetermined scale M . If $M \neq 0$ the spontaneous breaking of scale invariance will require the appearance of a Goldstone boson at the point $u = u_c$. The massless Goldstone boson is associated here with the spontaneous breaking of scale invariance [9]. Moreover, since the ground state is supersymmetric, we expect the appearance of a massless $O(N)$ singlet Goldstone boson (a dilaton) and its massless fermionic partner (a “dilantino”). In order to verify these properties, we now calculate the $\langle LL \rangle$ propagator that will enable us to see these poles in the appropriate four-point functions.

2.4 $\langle LL \rangle$ propagator, bound states and massless fermion and boson $O(N)$ singlet bound states

The $\langle LL \rangle$ propagator. We now calculate the $\langle LL \rangle$ propagator in the symmetric phase. Then, the propagators of the superfields ϕ and L are decoupled. In the example of a quartic potential, the ρ field can be eliminated by gaussian

integration. The relevant part of the L -action then reads

$$-\frac{N}{2u} \int d^3x d^2\theta (L - \mu)^2 + \frac{1}{2}(N-1) \text{Str} \ln (-\bar{\text{D}}\text{D} + 2L).$$

The calculation of the $\langle LL \rangle$ propagator involves the super-propagator (2.7). For the inverse propagator Δ_L^{-1} one finds

$$\Delta_L^{-1}(p) = -\frac{N}{u} \delta^2(\theta' - \theta) - 2N \int \frac{d^3k}{(2\pi)^3} \Delta(k, \theta, \theta') \Delta(p - k, \theta, \theta')$$

with, here, (see Eq. (A1.6))

$$\Delta(k, \theta, \theta') = \frac{1}{k^2 + M^2} [1 + \frac{1}{2}M\delta^2(\theta' - \theta)] e^{i\bar{\theta}\not{k}\theta'}.$$

Then,

$$\Delta(k, \theta, \theta') \Delta(p - k, \theta, \theta') = \frac{[1 + M\delta^2(\theta' - \theta)] e^{i\bar{\theta}\not{k}\theta'}}{(k^2 + M^2)[(p + k)^2 + M^2]}.$$

Notice the cancellation of the factor $e^{i\bar{\theta}\not{k}\theta'}$ which renders the integral more convergent than one could naively expect. The integral over k then yields the three-dimensional bubble diagram $B(p)$:

$$B(p) = \frac{1}{(2\pi)^3} \int \frac{d^3k}{(k^2 + M^2)[(p + k)^2 + M^2]} = \frac{1}{4\pi p} \text{Arctan}(p/2|M|).$$

At leading order for p small, we need only $B(0) = 1/8\pi|M|$. Then,

$$\Delta_L^{-1} = -\frac{N}{4\pi|M|} [1 + \{M + |M|(u_c/u)\}\delta^2(\theta' - \theta)] e^{i\bar{\theta}\not{p}\theta'}.$$

Considering that in momentum space we have

$$[-\bar{\text{D}}\text{D} + 2\mu]\delta^2(\theta' - \theta) = 2[-2 + \delta^2(\theta' - \theta)\mu] e^{-i\bar{\theta}\not{\mu}\theta'}, \quad (2.30)$$

we conclude that for small $M > 0$ the LL propagator corresponds to a super-particle of mass $2M(1 + u_c/u)$. For $M < 0$ the mass is $2|M(1 - u_c/u)|$. For $|u - u_c|$ small, it is a bound state and at the special point $u = u_c$ the mass vanishes.

More generally, we find

$$\Delta_L(p) = \frac{2}{NB(p)} \frac{1}{p^2 + m^2(p)} [1 - \frac{1}{2}m(p)\delta^2(\theta' - \theta)] e^{i\bar{\theta}\not{p}\theta'} \quad (2.31)$$

with

$$m(p) = 2M + \frac{1}{uB(p)}.$$

We note that the only renormalization required at leading order is a mass renormalization, a situation similar to the φ^6 scalar field theory [6]. As a consequence field dimensions are not modified.

Clearly, the propagation of the fields $M(x)$ and $\ell(x)$, as indicated in Eq. (2.31), when combined with the $L(\theta)\Phi^2$ interaction in Eq. (2.3), namely,

$$\int d^2\theta L\Phi^2 = M(-\tfrac{1}{2}\bar{\psi}\psi + F\varphi) - \varphi\bar{\ell}\psi + \tfrac{1}{2}\lambda\varphi^2,$$

describes the bound states in the $\varphi\varphi$, $\psi\psi$ and $\psi\varphi$ scattering amplitudes. For example [9], in the supersymmetric ground state case and with $\mu - \mu_c = 0$, the $\psi\varphi$ scattering amplitude $T_{\psi\varphi,\psi\varphi}(p^2)$, in the limit $p^2 \rightarrow 0$ satisfies

$$T_{\psi\varphi,\psi\varphi}(p^2) \sim \frac{2u}{N} \left[1 + \frac{u}{4\pi} \frac{M}{|M|} + \frac{u}{2\pi} \frac{i\not{p}}{|M|} \right]^{-1} \rightarrow -\frac{4\pi i}{N} \frac{|M|}{\not{p}} \quad (2.32)$$

for $M < 0$ and $u \rightarrow u_c$

One notes here that the fermion massless bound state pole appears when a non-zero solution (M) to the gap equation exists ($m_\varphi = m_\psi = |M|$) in the absence of any dimensional parameters ($\mu - \mu_c = 0$). This happens when the force between the massive ψ and φ quanta is determined by $u \rightarrow u_c$. The massless $O(N)$ singlet fermion bound state excitation is associated with the spontaneous breaking of scale invariance. Similarly, the bosonic partner of this massless bound state excitation can be then seen, at the same value of the parameters, in $\varphi\varphi$ and $\psi\psi$ scattering amplitudes as Eq. (2.31) shows.

At $\mu = \mu_c$ in the generic situation $M = 0$, or for $|p| \rightarrow \infty$ we find

$$B(p) = \frac{1}{8p}$$

and, thus,

$$\Delta_L^{-1}(p) = -\frac{N}{u}\delta^2(\theta' - \theta) - \frac{N}{4p}e^{i\bar{\theta}\not{p}\theta'}.$$

As a consequence the canonical dimension of the field L is one, as in perturbation theory, and the interaction $L\Phi^2$ in Eq. (2.3) is renormalizable.

2.5 Dimensions $2 \leq d \leq 3$

In d space-time dimensions $2 \leq d \leq 3$, the space of γ matrices implies $\text{tr } \mathbf{1} = 2$ and one notices, after some calculations following the same pattern as discussed above, that Eqs. (2.13) are replaced by

$$\begin{aligned} \rho - \varphi^2/N &= \Omega_d(m), \\ s - 2F\varphi/N &= 2M [\Omega_d(m) - \Omega_d(|M|)]. \end{aligned} \quad (2.33a)$$

Taking into account the other saddle point equations and $\rho_c = \Omega_d(0)$, one obtains a generalization of Eq. (2.19):

$$2\mathcal{E}/N = M^2\varphi^2/N + (M^2 - m^2) [\Omega_d(m) - \Omega_d(0)] + \int_0^\lambda d\lambda' \Omega_d(\sqrt{M^2 + \lambda'}) \quad (2.34)$$

and, thus,

$$\mathcal{E}/N = \frac{1}{2}M^2\varphi^2/N + K(d) \left[\frac{1}{2}(m^2 - M^2)m^{d-2} - (m^d - |M|^d)/d \right],$$

where

$$K(d) = -\frac{\Gamma(1 - d/2)}{(4\pi)^{d/2}}.$$

Moreover, from Eq. (2.34),

$$\frac{\partial \mathcal{E}}{\partial m} = -\frac{N}{2m}(m^2 - M^2)\Omega'_d(m) \quad (2.35)$$

has the sign of $m - |M|$ because $\Omega'_d(m)$ is negative. The function, therefore, has a minimum at $m = |M|$ for all values of d , and supersymmetry is maintained in the ground state for all $2 \leq d \leq 3$.

Two dimensions. It may be interesting to consider the same model in two dimensions. The model now is super-renormalizable. Even at $\mu = 0$ it is not chiral invariant since a chiral transformation that changes $\bar{\psi}\psi$ in $-\bar{\psi}\psi$, is equivalent to the change $U \mapsto -U$.

The expression (2.35) is cut-off independent and has a limit for $d = 2$:

$$\mathcal{E}/N = \frac{1}{2}M^2\varphi^2/N + \frac{1}{8\pi} [m^2 - M^2 - 2M^2 \ln(m/M)], \quad (2.36)$$

an expression which again has an absolute minimum at $m = |M|$. Then, a minimization with respect to M and φ yields $M\varphi = 0$. At the minimum $\mathcal{E}$ vanishes.

Taking into account $\lambda = 0$ and Eq. (2.33a) in the $d = 2$ limit, we obtain the gap equation

$$M = \mu + u\varphi^2/N + \frac{u}{2\pi} [\ln(\Lambda/|M|) + O(1)].$$

The solution $M^2 = \lambda = 0$, therefore, is not acceptable. The $O(N)$ symmetry is never broken. When the mass M is small in the cut-off scale, the l.h.s. is negligible and

$$m = M \propto \Lambda e^{2\pi\mu/u}.$$

This solution exists only when $-\mu/u$ is positive and large.

2.6 A supersymmetric non-linear σ -model at large N

We now consider the supersymmetric non-linear σ -model in d dimensions, $2 \leq d \leq 3$. The action

$$\mathcal{S}(\Phi) = \frac{1}{2\kappa} \int d^d x d^2 \theta \bar{D}\Phi \cdot D\Phi \quad (2.37)$$

involves a scalar superfield Φ , an N -component vector that satisfies

$$\Phi \cdot \Phi = N.$$

This relation is implemented by introducing a superfield

$$L(x, \theta) = M(x) + \bar{\theta}\ell(x) + \frac{1}{2}\bar{\theta}\theta\lambda(x),$$

where $M(x)$, $\lambda(x)$ and $\ell(x)$ are the Lagrange multiplier fields, and adding to the action

$$\mathcal{S}_L = \frac{1}{\kappa} \int d^d x d^2 \theta L(x, \theta) [\Phi^2(x, \theta) - N]. \quad (2.38)$$

The partition function is given by $(\mathcal{S}(\Phi, L) = \mathcal{S} + \mathcal{S}_L)$

$$\mathcal{Z} = \int [d\Phi][dL] e^{-\mathcal{S}(\Phi, L)}.$$

In terms of component fields, the total action reads

$$\begin{aligned} \mathcal{S} = \frac{1}{\kappa} \int d^d x \{ & \frac{1}{2}\varphi(-\partial^2 + M^2)\varphi - \frac{1}{2}\bar{\psi}(\not{\partial} + M)\psi \\ & + \frac{1}{2}\lambda(\varphi^2 - N) - \bar{\ell}(\bar{\psi} \cdot \varphi) \}. \end{aligned} \quad (2.39)$$

As in the case of the Φ^4 theory, we integrate out $N - 1$ superfield components leaving out $\phi = \Phi_1$:

$$\mathcal{Z} = \int [d\phi][dL] e^{-\mathcal{S}_N(\phi, L)},$$

where

$$\mathcal{S}_N(\phi, L) = \frac{1}{\kappa} \int d^d x d^2 \theta \left[\frac{1}{2} \bar{D}\phi \cdot D\phi + L(\phi^2 - N) \right] + \frac{N-1}{2} \text{Str} \ln(-\bar{D}D + 2L). \quad (2.40)$$

By varying the effective action with respect to ϕ , one finds the saddle point equation

$$\bar{D}D\phi - 2L\phi = 0, \quad (2.41)$$

which implies, for constant φ and $\psi = 0$,

$$F - M\varphi = 0 \quad \text{and} \quad MF + \lambda\varphi = 0. \quad (2.42)$$

Varying the large N action with respect to L and using the expression (2.7) for the ϕ propagator, one finds ($N \gg 1$)

$$\frac{1}{\kappa} - \frac{\phi^2}{N} = \int \frac{d^d k}{(2\pi)^d} \left[\frac{1 + M\bar{\theta}\theta}{k^2 + M^2 + \lambda} - \frac{M\bar{\theta}\theta}{k^2 + M^2} \right]. \quad (2.43)$$

We now introduce the boson mass parameter $m = \sqrt{M^2 + \lambda}$. In terms of its components Eq. (2.43) yields (for $\psi = 0$)

$$1 - \frac{\varphi^2}{N} = \kappa \Omega_d(m), \quad (2.44a)$$

$$M \frac{\varphi^2}{N\kappa} = M [\Omega_d(|M|) - \Omega_d(m)]. \quad (2.44b)$$

Dimension $d = 3$. Introducing the critical (cut-off dependent) value κ_c :

$$\frac{1}{\kappa_c} = \Omega_3(0),$$

we can write Eqs. (2.44) now as

$$\frac{1}{\kappa} - \frac{1}{\kappa_c} - \frac{\varphi^2}{N\kappa} = -\frac{m}{4\pi}, \quad (2.45a)$$

$$M \frac{\varphi^2}{N\kappa} = \frac{1}{4\pi} M (m - |M|). \quad (2.45b)$$

The calculation of the ground state energy density of the non-linear σ -model follows the similar steps as in the $(\Phi^2)^2$ model. One finds the action density

$$\mathcal{E}/N = \frac{1}{2N\kappa} m^2 \varphi^2 - \frac{1}{2} (m^2 - M^2) \left(\frac{1}{\kappa} - \frac{1}{\kappa_c} \right) - \frac{1}{12\pi} (m^3 - |M|^3) \quad (2.46)$$

and, taking into account the saddle point Eq. (2.44a),

$$\mathcal{E}/N = \frac{1}{2N\kappa} M^2 \varphi^2 + \frac{1}{24\pi} (m - |M|)^2 (m + 2|M|), \quad (2.47)$$

an expression identical to (2.19), up to the normalization of φ . Again, if a supersymmetric solution exists it has the lowest energy. We thus look for supersymmetric solutions.

In the $O(N)$ symmetric phase ($\varphi = 0$), Eq. (2.45b) is satisfied while Eq. (2.45a) yields

$$m = |M| = 4\pi \left(\frac{1}{\kappa_c} - \frac{1}{\kappa} \right).$$

This phase exists for $\kappa \geq \kappa_c$.

In the broken phase $m = M = 0$, and Eq. (2.45b) is again satisfied. Eq. (2.45a) then yields

$$\varphi^2/N = 1 - \kappa/\kappa_c, \quad (2.48)$$

which is the solution for $\kappa \leq \kappa_c$.

Since we have found solutions for all values of κ , we conclude that the ground state is always supersymmetric. At κ_c , a phase transition occurs between a massless phase with broken $O(N)$ symmetry ($\varphi^2 \neq 0$) and a symmetric phase with massive fermions and bosons of equal mass. The phase structure of the supersymmetric non-linear σ -model is analogous to the structure of the usual non-linear σ -model and less surprising than in the $(\Phi^2)^2$ field theory.

Dimension $d = 2$. The phase structure of the supersymmetric non-linear σ -model in two dimensions is rather simple, and again analogous to the structure of the usual non-linear σ -model. Eq. (2.45a) immediately implies that $M^2 + \lambda = 0$ is not a solution and thus the $O(N)$ symmetry remains unbroken ($\varphi^2 = 0$) for all values of the coupling constant κ . Then, with a suitable normalization of the cut-off Λ ,

$$m = \Lambda e^{-2\pi/\kappa}. \quad (2.49)$$

Correspondingly, the action density becomes

$$\frac{\mathcal{E}}{N} = -\frac{\lambda}{2\kappa} - \frac{1}{8\pi} [(M^2 + \lambda) \ln(M^2 + \lambda) - \lambda \ln \Lambda^2 - M^2 \ln M^2 - \lambda] \quad (2.50)$$

and, taking into account Eq. (2.49),

$$\frac{\mathcal{E}}{N} = \frac{1}{8\pi} [m^2 - M^2 + 2M^2 \ln(M/m)].$$

We recognize expression (2.36), and conclude in the same way that a supersymmetric solution has the lowest energy. Supersymmetry remains unbroken, and the common boson and fermion mass is

$$M = m = \Lambda e^{-2\pi/\kappa}. \quad (2.51)$$

The condition $M \ll \Lambda$ implies that the non-trivial physics is concentrated near $\kappa = 0$.

3 $O(N)$ supersymmetric model at finite temperature

We now study the phase structure of a supersymmetric model in $d = 3$ space-time dimensions at finite temperature.

3.1 The free energy at finite temperature

We start again from the action (2.1), which is transformed into

$$\mathcal{S}(\Phi, \rho, L) = \int d^3x d^2\theta \left[\frac{1}{2} \bar{D}\Phi \cdot D\Phi + NU(\Phi^2/N) + L(\theta)(\Phi^2(\theta) - NR(\theta)) \right],$$

$\Phi(\theta, x)$, $L(\theta, x)$ and $R(\theta, x)$ being N -component scalar superfields, parametrized as in the $T = 0$ discussion.

Integrating over $N - 1$ superfield components of Φ and keeping $\Phi_1 \equiv \phi$ (the scalar component of the superfield ϕ is identified as $\varphi_1 \equiv \varphi$) one finds an expression for the partition function $\mathcal{Z}$ that differs from Eq. (2.4) only by the field boundary conditions

$$\mathcal{Z} = \int [d\phi][dR][dL] e^{-\mathcal{S}_N(\phi, R, L)}, \quad (3.1)$$

where $\mathcal{S}_N$ is the large N action (2.5):

$$\begin{aligned} \mathcal{S}_N = \int d^3x d^2\theta & \left[\frac{1}{2} \bar{D}\phi D\phi + NU(R) + L(\phi^2 - NR) \right] \\ & + \frac{1}{2}(N-1) \text{Str} \ln [-\bar{D}D + 2L]. \end{aligned} \quad (3.2)$$

Among the three saddle point equations at $T = 0$, only one is changed at finite temperature:

$$2L\phi - \bar{D}D\phi = 0, \quad (3.3a)$$

$$L - U'(R) = 0, \quad (3.3b)$$

$$R - \phi^2/N = \frac{1}{N} \text{tr} \Delta(k, \theta, \theta). \quad (3.3c)$$

Eq. (3.3c) involves $\Delta(k, \theta, \theta)$ (Eq. (2.8))

$$\Delta(k, \theta, \theta) = \frac{1}{k^2 + M_T^2 + \lambda} + \bar{\theta}\theta M_T \left(\frac{1}{k^2 + M_T^2 + \lambda} - \frac{1}{k^2 + M_T^2} \right), \quad (3.4)$$

where we call M_T the expectation value of $M(x)$ at finite temperature T .

When written in components, Eq. (3.3a) implies

$$F - M_T \varphi = 0, \quad (3.5a)$$

$$\lambda \varphi + M_T F = 0. \quad (3.5b)$$

and, thus, the Goldstone condition $\varphi(\lambda + M_T^2) = 0$ follows.

Eq. (3.3b) implies

$$M_T = U'(\rho), \quad \lambda = sU''(\rho). \quad (3.6)$$

When calculating the trace in Eq. (3.3c), we have to take into account that bosons at finite temperature satisfy periodic and fermions anti-periodic boundary conditions. For bosons we introduce the function

$$\begin{aligned} G_2(m_T, T) &= \frac{T}{(2\pi)^{d-1}} \sum_{n \in \mathbb{Z}} \int^\Lambda \frac{d^{d-1}k}{(2\pi nT)^2 + k^2 + m_T^2} \\ &= \int^\Lambda \frac{d^{d-1}k}{(2\pi)^{d-1}} \frac{1}{\omega(k)} \left(\frac{1}{2} + \frac{1}{e^{\beta\omega_\varphi(k)} - 1} \right). \end{aligned} \quad (3.7)$$

with $\omega_\varphi(k) = \sqrt{M_T^2 + \lambda + k^2}$. It is convenient here to introduce the boson thermal mass

$$m_T = \sqrt{M_T^2 + \lambda}. \quad (3.8)$$

The parameter m_T characterizes the decay of correlation functions in space directions. Note that M_T , however, does not characterize the decay of fermion correlations, because fermions have no zero mode, the relevant parameter being $\sqrt{M_T^2 + \pi^2 T^2}$.

Combining Eq. (3.7) with the $\theta = 0$ part of the finite temperature Eq. (3.3c) we obtain

$$\begin{aligned} \rho - \varphi^2/N &= \int \frac{d^2k}{(2\pi)^2} \frac{1}{\omega_\varphi(k)} \left(\frac{1}{2} + \frac{1}{e^{\omega_\varphi(k)/T} - 1} \right) \\ &= \rho_c - \frac{T}{2\pi} \ln(2 \sinh(m_T/2T)), \end{aligned} \quad (3.9)$$

where ρ_c has been defined in Eq.(2.15).

In the same manner, for fermions we introduce the function

$$\mathcal{G}_2(M_T, T) = \int^\Lambda \frac{d^{d-1}k}{(2\pi)^{d-1}} \frac{1}{\omega(k)} \left(\frac{1}{2} - \frac{1}{e^{\omega_\psi(k)/T} + 1} \right) \quad (3.10)$$

with $\omega_\psi(k) = \sqrt{k^2 + M_T^2}$. (This is the fermion analogue of Eq. (3.7) in which one recognizes again the sum of quantum and thermal contributions. Note that the function $\mathcal{G}_2(M_T, T)$ has a regular expansion in M_T^2 at $M_T = 0$.) Combined with the $\bar{\theta}\theta$ part of Eq. (3.3c), it leads to

$$\begin{aligned} s - 2F\varphi/N &= 2M_T \int \frac{d^2k}{(2\pi)^2} \left\{ \frac{1}{2\omega_\varphi(k)} \left(\frac{1}{2} + \frac{1}{e^{\omega_\varphi(k)/T} - 1} \right) \right. \\ &\quad \left. - \frac{1}{2\omega_\psi(k)} \left(\frac{1}{2} - \frac{1}{e^{\omega_\psi(k)/T} + 1} \right) \right\}. \end{aligned}$$

Integrating we obtain

$$s - 2F\varphi/N = \frac{TM_T}{\pi} [\ln(2 \cosh(M_T/2)) - \ln(2 \sinh(m_T/2T))] . \quad (3.11)$$

The action density $\mathcal{F}$ at finite temperature, to leading order in $1/N$, is given by ($\beta = 1/T$)

$$\mathcal{F} = \mathcal{S}_N/V_2\beta , \quad (3.12)$$

where $\mathcal{S}_N$ is given by Eq. (3.2) at constant fields and V_2 is the two-dimensional volume.

The supertrace term in Eq. (3.2) decomposes into boson and fermion contributions. The boson contribution involves ($\omega_n = 2n\pi T$):

$$\begin{aligned} \frac{1}{V_{d-1}} \text{tr} \ln(-\nabla^2 + m_T^2) &= \int \frac{d^{d-1}k}{(2\pi)^{d-1}} \sum_{n \in \mathbb{Z}} \ln(\omega_n^2 + k^2 + m_T^2) \\ &= 2 \int \frac{d^{d-1}k}{(2\pi)^{d-1}} \ln [2 \sinh(\beta\omega_\varphi(k)/2)] , \end{aligned} \quad (3.13)$$

where V_{d-1} is the $(d-1)$ -dimensional volume. The result in Eq. (3.13) is given up to a mass independent infinite constant.

Similarly, for fermions (now $\omega_n = (2n+1)\pi T$)

$$\begin{aligned} \frac{1}{V_{d-1} \text{tr} \mathbf{1}} \text{tr} \ln(\not{\partial} + M_T) &= \frac{1}{2} \int \frac{d^{d-1}k}{(2\pi)^{d-1}} \sum_{n \in \mathbb{Z}} \ln(\omega_n^2 + k^2 + M_T^2) \\ &= \int \frac{d^{d-1}k}{(2\pi)^{d-1}} \ln [2 \cosh(\omega_\psi(k)/2T)] . \end{aligned} \quad (3.14)$$

Thus, one finds

$$\begin{aligned} \frac{1}{V_2} \text{Str} \ln [-\bar{\text{D}}\text{D} + 2L] &= \frac{1}{V_2} \text{tr} \ln(-\partial^2 + M_T^2 + \lambda) - \frac{1}{V_2} \text{tr} \ln(\not{\partial} + M_T) \\ &= 2 \int \frac{d^2k}{(2\pi)^2} \ln[2 \sinh(\beta\omega_\varphi/2)] - 2 \int \frac{d^2k}{(2\pi)^2} \ln[2 \cosh(\beta\omega_\psi/2)] \\ &= \frac{1}{T} \rho_c(m_T^2 - M_T^2) - \frac{1}{6\pi T} (m_T^3 - |M_T|^3) \\ &\quad + 2 \int \frac{d^2k}{(2\pi)^2} \{ \ln[1 - e^{-\beta\omega_\varphi}] - \ln[1 + e^{-\beta\omega_\psi}] \} . \end{aligned} \quad (3.15)$$

The explicitly subtracted part reduces for $M_T = M$ to the $T = 0$ result. The other contributions to $\mathcal{F}$ are the same (up to the change $M \mapsto M_T$) as in Eq. (2.17) and, therefore, one finds

$$\begin{aligned} \frac{1}{N} \mathcal{F} &= -\frac{F^2}{2N} + M_T \frac{F\varphi}{N} + \lambda \frac{\varphi^2}{2N} + \frac{1}{2} s(U'(\rho) - M_T) - \frac{1}{12\pi} (m_T^3 - |M_T|^3) \\ &\quad + \frac{1}{2} \lambda (\rho_c - \rho) + T \int \frac{d^2k}{(2\pi)^2} \{ \ln[1 - e^{-\beta\omega_\varphi}] - \ln[1 + e^{-\beta\omega_\psi}] \} , \end{aligned} \quad (3.16)$$

which reduces at $T = 0$ to the action density of Eq. (2.17), $\mathcal{F}(T = 0) \equiv \mathcal{E}$.

Eq. (3.16) can be rewritten by using Eq. (3.9), which is also obtained by setting to zero the $\partial/\partial\lambda$ derivative of Eq. (3.16), as well as $M_T - U'(\rho) = 0$ from Eq. (3.6) (or equivalently, setting to zero the $\partial/\partial s$ derivative of Eq. (3.16)), one finds

$$\begin{aligned} \frac{1}{N}\mathcal{F} = & \frac{1}{2}M_T^2\frac{\varphi^2}{N} + \frac{1}{24\pi}(m_T - |M_T|)^2(m_T + 2|M_T|) \\ & + \frac{T\lambda}{4\pi}\ln(1 - e^{-m_T/T}) + T \int \frac{d^2k}{(2\pi)^2} \{\ln(1 - e^{-\beta\omega_\varphi}) - \ln(1 + e^{-\beta\omega_\psi})\}. \end{aligned} \quad (3.17)$$

Eq. (3.17) is the finite temperature version of Eq. (2.19).

Inserting Eqs. (3.9) and (3.11) (with $F = M_T\varphi$) into Eqs. (3.6) with $U(\rho) = \mu\rho + \frac{1}{2}u\rho^2$ one finds ($\mu_c = -u\rho_c$)

$$M_T = \mu - \mu_c + u\frac{\varphi^2}{N} - \frac{u}{2\pi}T \ln\left(2 \sinh\left(\frac{1}{2}m_T/T\right)\right), \quad (3.18)$$

$$\frac{m_T^2 - M_T^2}{uM_T} = \frac{2\varphi^2}{N} - \frac{T}{\pi} \left[\ln\left(2 \sinh\left(\frac{1}{2}m_T/T\right)\right) - \ln\left(2 \cosh\left(\frac{1}{2}|M_T|/T\right)\right) \right], \quad (3.19)$$

which are the gap equations for $T \neq 0$.

Solution to the saddle point equations: We first note that the r.h.s. of the gap Eq. (3.18) diverges when $m_T \rightarrow 0$. The finite temperature system in three dimensions has the property of a statistical system in two dimensions. In two dimensions a spontaneous breaking of a continuous symmetry is impossible, due to the IR behaviour of a system with potential massless Goldstone particles. Therefore, the $O(N)$ symmetry is never broken, $\varphi = 0$, and thus,

$$\frac{M_T}{T} = \frac{\mu - \mu_c}{T} - \frac{u}{2\pi} \ln\left(2 \sinh\left(\frac{1}{2}m_T/T\right)\right), \quad (3.21)a$$

$$\frac{M_T^2 - m_T^2}{uTM_T} = \frac{1}{\pi} \left[\ln\left(2 \sinh\left(\frac{1}{2}m_T/T\right)\right) - \ln\left(2 \cosh\left(\frac{1}{2}M_T/T\right)\right) \right]. \quad (3.22)b$$

Note that while the expression of $\mathcal{F}$ is complicated, its derivative with respect to m_T^2 remains simple:

$$\frac{1}{N} \frac{\partial \mathcal{F}}{\partial m_T^2} = \frac{(m_T^2 - M_T^2)}{16\pi m_T \tanh(m_T/2T)}. \quad (3.23)$$

Therefore, the minimum still occurs at $m_T = |M_T|$, but one verifies that $m_T = |M_T|$ is not a solution to the saddle point equations. We have only found a lower bound on the free energy density $\mathcal{F}$ as a function of M_T :

$$\mathcal{F} = NT \int \frac{d^2k}{(2\pi)^2} \ln \tanh\left(\frac{1}{2}\beta\sqrt{M_T^2 + k^2}\right). \quad (3.24)$$

Its derivative with respect to M_T is

$$\frac{\partial \mathcal{F}}{\partial M_T} = -N \frac{T M_T}{2\pi} \ln \tanh(|M_T|/2T).$$

Therefore, the derivative has the sign of M_T . The lower bound has a limit for $M_T \rightarrow 0$, which is thus an absolute lower bound:

$$\mathcal{F} = -\frac{7}{8\pi} \zeta(3) N T^3. \quad (3.25)$$

Similarly, the derivative of $\mathcal{F}$ with respect to M_T at m_T fixed is

$$\begin{aligned} \frac{\partial \mathcal{F}}{\partial M_T} &= \frac{M_T}{2\pi} [\ln(2 \cosh(M_T/2T)) - \ln(2 \sinh(m_T/2T))] \\ &= \frac{m_T^2 - M_T^2}{2u}. \end{aligned}$$

We now have to find the solutions to the saddle point equations and compare their free energies.

Note a first solution in the regime $T \rightarrow 0$, $\mu < \mu_c$ with $|m_T| \ll T$ and $|M_T| \ll T$. Then, from (3.18) we find the boson thermal mass

$$m_T \sim T e^{-2\pi(\mu_c - \mu)/uT}.$$

Eq. (3.19) yields the fermion thermal mass

$$M_T \sim \frac{m_T^2}{\mu_c - \mu}.$$

Since both m_T and M_T are very small, the free energy is very close to the lower bound (3.25). The solution we have found here is a precursor of the zero-temperature phase transition, and corresponds to $\varphi \neq 0$ in region III and IV of Fig. 1.

Other solutions exist in this regime but they converge, up to exponential corrections, to the finite masses of the $T = 0$ spectrum and, thus, their free energy is much larger from the lower bound (3.24).

It is expected that even for T larger and $\mu \geq \mu_c$, the continuation of the small mass solution remains the solution with lowest energy.

In the following we take $m \equiv m_T$ as the free variable and, using Eq. (3.18) for $M_T = M_T(m, \varphi, \mu, u, T)$, we discuss

$$W(m, \varphi, T) = \frac{1}{N} \mathcal{F}(m_T = m, M_T(m, \varphi, \mu, u, T), T). \quad (3.26)$$

Since, as described above, at $T \neq 0$ there is no breaking of the $O(N)$ symmetry, we will discuss mainly $\varphi = 0$ in Eq. (3.26). For $T = 0$, Eq. (3.26) results in $\frac{1}{N}\mathcal{F}(m, \varphi^2, T = 0) = \frac{1}{N}\mathcal{E}(m, \varphi^2)$ which has been analyzed above. In particular we see the phase structure in Fig.1 and the interesting degeneracy found in Regions II and IV and exhibited in Figs. 3-4. Rather than changing the value of $\mu - \mu_c$ and the coupling u as has been done above at $T = 0$, we will be interested here to see the temperature dependence while these parameters will be held fixed.

At finite temperature, the effective field theory describes the interactions of the fermions and bosons with the heat bath. This interaction acts like a source of soft breaking of supersymmetry. The short distance behavior is cured at finite temperature in a similar way it happens at $T = 0$. We now show that the general properties of the transitions as a function of the temperature T have a similar character as the transitions seen at $T = 0$ when the coupling constant is varied.

As seen in Fig. 1, at $T = 0$ there are two regions in the $\{\mu - \mu_c, u\}$ space where the vacua are degenerate. These are (region **II** : $\mu - \mu_c \geq 0$ and $u \geq u_c$ and region **IV** : $\mu - \mu_c \leq 0$ and $u \leq u_c$). Fig. 2 shows the ground state free energy at $T = 0$ in region **IV**.

We discuss first region (**II**) $\mu - \mu_c \geq 0$ and $u \geq u_c$:

Here, as T increases the ground state with the smaller mass ($m_\varphi \equiv m = m_+$) has a lower free energy than the heavier one ($m_\varphi \equiv m = m_- > m_+$) due to a sizable entropy contribution. (Fig. 5).

Peculiar transitions can occur in this system. If the system was initially at $T = 0$ in the ground state with a boson mass $m_\varphi = m_-$, it will eventually go, as T is increased, into the state with mass $m_\varphi = m_+$, namely, into the lower mass ground state. This is due to the entropy negative contribution to the free energy forcing the system to favors the lowest mass state. On the other hand, a system that started at a high temperature in the state of low mass $m_\varphi = m_+$ will stay in this state as the system is cooled and will never roll back into the $m = m_-$ high mass phase. Favoring the lowest mass phase as the temperature increases is a general effect that will occur in any physical system that is initially (at $T = 0$) mass degenerated. Here, supersymmetry imply that the m_+ and m_- vacua are at the same energy $E = 0$ at $T = 0$.

Region (**IV**) $\mu \leq 0$ and $u \leq u_c$: here one finds at $T = 0$ two distinct degenerate phases. One is an ordered phase ($\varphi \neq 0$) with a zero boson and fermion mass, the other is a symmetric phase ($\varphi = 0$) with a massive ($m = m_-$) boson and fermion (as seen in Fig. 2).

As the temperature increases, the ordered phase with the broken $O(N)$ symmetry ($m = 0, \varphi^2 \neq 0$) disappears into a $\varphi^2 = 0$ symmetric phase and a very small mass ground state $m_\varphi = m \geq 0$ is created (as seen in Fig. 6 and in Fig. 7).

Eventually, also the other $O(N)$ symmetric vacuum ($m = m_-, \varphi^2 = 0$) goes into the small mass $O(N)$ -symmetric ground state.

As in **II**, a system that was initially, at $T = 0$, in the $O(N)$ -symmetric phase




Fig. 5 The ground state free energy $\frac{1}{N}\mathcal{F}(m \equiv m_\varphi, \varphi, T)$ as given in Eq. (3.17) at $\varphi = 0$ as a function of the boson mass (m) at different temperatures. Here $\mu - \mu_c = 1$ (this sets the mass scale), $u/u_c = 1.5$ and T varies between $T = 0 - -0.5$. At $T = 0$ two degenerate $O(N)$ symmetric phases exist with a light $m = m_+$ and heavier $m = m_- > m_+$ massive boson (and fermion). As the temperature increases, the light mass phase is stronger affected as its entropy increases faster and becomes the only ground state.

$(m_\varphi = m_-, \varphi^2 = 0)$ will decay into the smaller mass state when the temperature is raised. But upon cooling the system in the small mass phase will roll into the ordered phase $(m_\varphi = m = 0, \varphi^2 \neq 0)$ at $T = 0$. The system will never roll back into the symmetric phase $(m_\varphi = m_-, \varphi = 0)$.

4 The supersymmetric $O(N)$ non-linear σ -model at finite temperature

The supersymmetric non-linear σ -model we consider here has already been discussed above at zero temperature. The partition function of the $O(N)$ supersymmetric non-linear σ -model in d dimensions is given by

$$\mathcal{Z} = \int [d\Phi][dL] e^{-\mathcal{S}(\Phi, L)} \quad (4.1)$$

with

$$\mathcal{S}(\Phi) = \frac{1}{2\kappa} \int d^d x d^2 \theta \bar{D}\Phi \cdot D\Phi + L(\Phi^2 - N), \quad (4.2)$$

where κ is a constant. The scalar superfield $\Phi(x, \theta)$ is an N -component vector and the scalar superfield $L(x, \theta)$ is parameterized as in the $T = 0$ case. After integrating out $N - 1$ components of Φ , leaving out $\phi = \Phi_1$, we obtain

$$\mathcal{Z} = \int [d\phi][dL] e^{-\mathcal{S}_N(\phi, L)}$$




Fig. 6 Same as Fig. 2 but the temperature has been increased from $T = 0$ (in Fig. 2) to $T = 0.7$ (here). $\frac{1}{N}\mathcal{F}(m \equiv m_\varphi, \varphi, T)$ as given in Eq. (3.17) as a function of the boson mass (m) and A , where $A^2 = \varphi^2/u_c$. Here $\mu - \mu_c = -1$ and $u/u_c = 0.2$. A non-degenerate $O(N)$ symmetric ground state ($\varphi = 0$) appears with a very small boson mass (the non-zero mass is not seen here due to the limited resolution of the plot. See Fig. 7).

where

$$\mathcal{S}_N(\phi, L) = \frac{1}{\kappa} \int d^d x d^2 \theta \left[\frac{1}{2} \bar{D}\phi \cdot D\phi + L(\phi^2 - N) \right] + \frac{N-1}{2} \text{Str} \ln(-\bar{D}D + 2L). \quad (4.3)$$

Differentiating the action $\mathcal{S}_N(\phi, L)$ with respect to the superfields ϕ and L , one obtains Eq. (2.41) and the generalization of Eq. (2.43):

$$2L\phi - \bar{D}D\phi = 0, \quad (4.4a)$$

$$\frac{N}{\kappa} - \frac{\phi^2}{N} = \text{tr} \Delta(k, \theta, \theta). \quad (4.4b)$$

The first equation is equivalent to $F = M_T \varphi$ and $\lambda \varphi + F M_T = 0$ while the second equation can be compared with Eq. (3.3c), and thus leads to (for $N \gg 1$ and $\psi = 0$)

$$\begin{aligned} \frac{1}{\kappa} - \frac{\varphi^2}{N} &= \int \frac{d^d k}{(2\pi)^d} \frac{1}{\omega_\varphi(k)} \left(\frac{1}{2} + \frac{1}{e^{\omega_\varphi(k)/T} - 1} \right), \\ -\frac{2F\varphi}{N} &= 2M_T \int \frac{d^d k}{(2\pi)^d} \left\{ \frac{1}{2\omega_\varphi(k)} \left(\frac{1}{2} + \frac{1}{e^{\omega_\varphi(k)/T} - 1} \right) \right\} \end{aligned} \quad (4.5a)$$




Fig. 7 This figure displays the effect of increasing the temperature from $T = 0$ in Fig. 2 to T that varies between $T = 0 - 0.7$ (Fig. 6 has $T = 0.7$). The ground state free energy $\frac{1}{N}\mathcal{F}(m \equiv m_\varphi, \varphi, T)$ as given in Eq. (3.17) at $\varphi = 0$ is plotted as a function of the boson mass (m) at different temperatures. Here $\mu - \mu_c = -1$, $u/u_c = 0.2$. At $T = 0$ there are two degenerate phases: An $O(N)$ -symmetric phase, shown here, with a massive ($m = m_-$) boson and fermion and an ordered phase ($\varphi \neq 0$) with massless particles (both phases are shown in Fig. 2). At finite temperatures the $O(N)$ symmetry is restored (see Fig. 6) and a small mass ground state appears, the heavy mass state decays into the small mass ground state as seen here.

$$-\frac{1}{2\omega_\psi(k)} \left(\frac{1}{2} - \frac{1}{e^{\omega_\psi(k)/T} + 1} \right) \} \quad (4.5b)$$

with $\omega_\varphi(k) = \sqrt{m_T^2 + k^2}$ and $\omega_\psi(k) = \sqrt{M_T^2 + k^2}$.

Dimension $d = 3$. As we have already discussed, the $d = 3$ finite temperature theory is analogous from the point of phase transitions to a two-dimensional theory. Therefore, the $O(N)$ symmetry remains unbroken and $\varphi = 0$. We thus write the two gap equations only in this limit:

$$\frac{1}{\kappa_c} - \frac{1}{\kappa} = \frac{T}{2\pi} \ln(2 \sinh(m_T/2T)), \quad (4.6a)$$

$$0 = \frac{TM_T}{2\pi} [\ln(2 \sinh(m_T/2T)) - \ln(2 \cosh(M_T/2T))]. \quad (4.6b)$$

The calculation of the ground state energy density to leading order for $N \rightarrow \infty$ of the non-linear σ -model at finite temperature follows steps similar to those of

the $(\Phi^2)^2$ field theory. The free energy at finite temperature is given by

$$\mathcal{F} = T\mathcal{S}_N/V_2, \quad (4.7)$$

where $\mathcal{S}_N$ is given by Eq. (4.3) at constant fields and V_2 is the two-dimensional volume. Here, the supertrace term in Eq. (4.3) has to be replaced by the finite temperature form as it appears in Eq. (3.16). The free energy then is given by

$$\begin{aligned} \frac{1}{N}\mathcal{F} = & \frac{1}{2}(M_T^2 - m_T^2) \left(\frac{1}{\kappa} - \frac{1}{\kappa_c} \right) - \frac{1}{12\pi} (m_T^3 - |M_T|^3) \\ & + T \int \frac{d^2k}{(2\pi)^2} \{ \ln[1 - e^{-\beta\omega_\varphi}] - \ln[1 + e^{-\beta\omega_\psi}] \}. \end{aligned} \quad (4.8)$$

The free energy in Eq. (4.8) can be compared with Eq. (2.46) which gives the ground state energy $\mathcal{F}$ at $T = 0$.

It can be verified that, after using the zero temperature limit of Eq. (4.6a), the energy density becomes identical to the energy density of the $(\Phi^2)^2$ theory, up to a possible rescaling of the field. It is now clear that the same mechanism works here. Using Eq. (4.6a), one indeed finds the expression (3.17) (for $\varphi = 0$):

$$\begin{aligned} \frac{1}{N}\mathcal{F} = & \frac{1}{24\pi} (m_T - |M_T|)^2 (m_T + 2|M_T|) + \frac{T}{4\pi} (m_T^2 - M_T^2) \ln(1 - e^{-m_T/T}) \\ & + T \int \frac{d^2k}{(2\pi)^2} \{ \ln(1 - e^{-\beta\omega_\varphi}) - \ln(1 + e^{-\beta\omega_\psi}) \}. \end{aligned} \quad (4.9)$$

Solutions. The derivative of $\mathcal{F}$ with respect to $|M_T|$ at m_T fixed is

$$\frac{\partial \mathcal{F}}{\partial |M_T|} = \frac{|M_T|T}{2\pi} [\ln(2 \cosh(M_T/2T)) - \ln(2 \sinh(m_T/2T))].$$

It is convenient to introduce the notation

$$X(\kappa, T) = \exp \left[\frac{2\pi}{T} \left(\frac{1}{\kappa_c} - \frac{1}{\kappa} \right) \right].$$

Then, the behaviour of $\mathcal{F}$ depends on the position of X with respect to 2:

$$X = 2 \Leftrightarrow T = \frac{2\pi}{\ln 2} \left(\frac{1}{\kappa_c} - \frac{1}{\kappa} \right),$$

an equation which has a solution only for $\kappa > \kappa_c$.

For $X < 2$, the derivative vanishes at $M_T = 0$ and is positive for all $|M_T| > 0$. Instead for $X > 2$, the derivative vanishes both at $M_T = 0$ and

$$|M_T| = 2T \ln \left[\frac{1}{2} (X + \sqrt{X^2 - 4}) \right], \quad (4.10)$$

which are the two solutions of Eq. (4.6a). The minimum of $\mathcal{F}$ is located at the second value (4.10).

We find, therefore, an interesting non-analytic behaviour:

$$\begin{cases} M_T = 0 & \text{for } X < 2, \\ M_T = 2T \ln \left[\frac{1}{2}(X + \sqrt{X^2 - 4}) \right] & \text{for } X > 2. \end{cases}$$

Eq. (4.6a) then yields directly the boson thermal mass

$$m_T = 2T \ln \left[\frac{1}{2}(X + \sqrt{X^2 + 4}) \right].$$

For $\kappa < \kappa_c$ and $T \rightarrow 0$, we find the asymptotic behaviour

$$m_T \sim TX(\kappa, T),$$

which is exponentially small, and $M_T = 0$.

For $\kappa > \kappa_c$ and $T \rightarrow 0$, both m_T and M_T converge toward the finite $T = 0$ limit with exponential corrections.

In the opposite high temperature limit $T \gg |1/\kappa - 1/\kappa_c|$, we find that m_T is asymptotically proportional to T :

$$m_T \sim 2T \ln((1 + \sqrt{5})/2),$$

while $M_T = 0$.

It is not clear whether such a result can survive $1/N$ corrections.

Dimension $d = 2$. In generic dimensions $2 \leq d \leq 3$, a phase transition is not even possible at finite temperature and $\varphi = 0$ (in two dimensions it is even impossible at zero temperature). The gap equations take the form

$$\frac{1}{\kappa} = \Omega_d(m_T) + T^{d-2} f_d(m_T^2/T^2), \quad (4.11)$$

$$0 = M_T [\Omega_d(|M_T|) - T^{d-2} g_d(M_T^2/T^2) - \Omega_d(m) - T^{d-2} f_d(m_T^2/T^2)], \quad (4.12)$$

where f_d and g_d are defined by

$$\begin{aligned} f_d(z) &= N_{d-1} \int_0^\infty \frac{x^{d-2} dx}{\sqrt{x^2 + z}} \frac{1}{\exp[\sqrt{x^2 + z}] - 1} \\ &= N_{d-1} \int_{\sqrt{z}}^\infty (y^2 - z)^{(d-3)/2} \frac{dy}{e^y - 1} \end{aligned} \quad (4.13)$$

and

$$\begin{aligned} g_d(s) &= N_{d-1} \int_0^\infty \frac{x^{d-2} dx}{\sqrt{x^2 + s}} \frac{1}{\exp[\sqrt{x^2 + s}] + 1} \\ &= N_{d-1} \int_{\sqrt{s}}^\infty (y^2 - s)^{(d-3)/2} \frac{dy}{e^y + 1}. \end{aligned} \quad (4.14)$$

For $d = 2$, it is then convenient to introduce the physical mass m solution of

$$1/\kappa = \Omega_2(m),$$

so that the equations become

$$\Omega_2(m) = \Omega_2(m_T) + f_2(m_T^2/T^2), \quad (4.15)$$

and either $M_T = 0$ or

$$\Omega_2(|M_T|) - g_2(M_T^2/T^2) - \Omega_2(m_T) - f_2(m_T^2/T^2) = 0. \quad (4.16)$$

The first equation which determines m_T is identical to the equation one obtains for the usual non-linear σ -model. At low temperature $m_T = m$ up to exponentially small corrections. At high temperature

$$\frac{T}{m_T} \sim \frac{1}{\pi} \ln(m_T/m) \sim \frac{1}{\pi} \ln(T/m),$$

a consequence of the domination of the zero mode and the UV asymptotic freedom of the non-linear σ -model.

Combining both gap equations we find

$$\Omega_2(|M_T|) - g_2(M_T^2/T^2) = \Omega_2(m).$$

The analysis of the equation for $d = 2$ shows that, for T large, the equation has no solution and thus $M_T = 0$, but it has a solution for T small. The situation, therefore, is similar to what has been encountered in three dimensions. Again, an analysis of LL propagator and $1/N$ corrections is required to understand whether this result survives beyond the large N limit.

Acknowledgements

MM would like to thank W.A. Bardeen and J. Feinberg for useful discussions. The warm hospitality of the Saclay and Fermilab theory groups is also appreciated.

Appendices

A1 Conventions: Supersymmetry and Majorana spinors in $d = 3$

We briefly recall the properties of Majorana spinors in three euclidean space dimensions and explain our notation. In three dimensions the spin group is $SU(2)$. Then, a spinor transforms like

$$\psi_U = U\psi, \quad U \in SU(2) \quad .$$

The role of Dirac γ matrices is played by the Pauli σ matrices, $\gamma_\mu \equiv \sigma_\mu$. Moreover, σ_2 is antisymmetric while $\sigma_2\sigma_\mu$ is symmetric. This implies

$$\sigma_2\sigma_\mu\sigma_2 = -{}^T\sigma_\mu \Rightarrow U^* = \sigma_2 U \sigma_2 \quad .$$

A Majorana spinor corresponds to a neutral fermion and has only two independent components ψ_1, ψ_2 .

The conjugated spinor is defined by (T means transposed)

$$\bar{\psi} = {}^T\psi\sigma_2 \Leftrightarrow \bar{\psi}_\alpha = i\epsilon_{\alpha\beta}\psi_\beta, \quad (A1.1)$$

($\epsilon_{\alpha\beta} = -\epsilon_{\beta\alpha}$, $\epsilon_{12} = 1$) and thus $\bar{\psi}$ transforms like

$$[{}^T\psi\sigma_2]_U = [{}^T\psi\sigma_2]U^\dagger.$$

One also defines a spinor of Grassmann coordinates:

$$\bar{\theta} = {}^T\theta\sigma_2 \quad ,$$

Since the only non-vanishing product is $\theta_1\theta_2$, we have

$$\bar{\theta}_\alpha\theta_\beta = \frac{1}{2}\delta_{\alpha\beta}\bar{\theta} \cdot \theta \quad .$$

The scalar product $\bar{\theta} \cdot \theta$ is

$$\bar{\theta} \cdot \theta = -2i\theta_1\theta_2 \Rightarrow \theta_\alpha\bar{\theta}_\beta = i\delta_{\alpha\beta}\theta_1\theta_2.$$

If $\bar{\theta}', \theta'$ is another pair of coordinates, because $\sigma_2\sigma_\mu$ is symmetric, one finds

$$\bar{\theta}'\sigma_\mu\theta' = {}^T\theta'\sigma_2\sigma_\mu\theta' = -\bar{\theta}'\sigma_\mu\theta, \quad (A1.2)$$

and for the same reason

$$\bar{\theta}\psi = \bar{\psi}\theta.$$

Other useful identities are

$$(\bar{\theta}\psi)^2 = -\frac{1}{2}(\bar{\theta}\theta)(\bar{\psi}\psi), \quad (\bar{\theta}\not{p}\psi)^2 = \frac{1}{2}p^2(\bar{\psi}\psi)(\bar{\theta}\theta).$$

It is convenient to integrate over θ_1, θ_2 with the measure

$$d^2\theta \equiv \frac{i}{2}d\theta_2d\theta_1.$$

Then,

$$\int d^2\theta \bar{\theta}_\alpha \theta_\beta = \frac{1}{2}\delta_{\alpha\beta}, \quad \int d^2\theta \bar{\theta} \cdot \theta = 1.$$

With this convention the identity kernel $\delta^2(\theta' - \theta)$ in θ space is

$$\delta^2(\theta' - \theta) = (\bar{\theta}' - \bar{\theta}) \cdot (\theta' - \theta). \quad (A1.3)$$

Superfields and covariant derivatives. A superfield $\Phi(\theta)$ has the expansion

$$\Phi(\theta) = \varphi + \bar{\theta}\psi + \frac{1}{2}\bar{\theta}\theta F. \quad (A1.4)$$

Again, although only two θ variables are independent, we define the covariant derivatives D_α and $\bar{D}_\alpha$ ($\bar{D} = \sigma_2 D$):

$$D_\alpha \equiv \frac{\partial}{\partial \bar{\theta}_\alpha} - (\not{\theta})_\alpha, \\ \bar{D}_\alpha \equiv \frac{\partial}{\partial \theta_\alpha} - (\bar{\theta}\not{\theta})_\alpha.$$

Then the anticommutation relation is

$$\{D_\alpha, \bar{D}_\beta\} = -2[\not{\theta}]_{\alpha\beta}.$$

Also

$$\bar{D}_\alpha D_\alpha = \frac{\partial}{\partial \theta_\alpha} \frac{\partial}{\partial \bar{\theta}_\alpha} - (\bar{\theta}\not{\theta})_\alpha \frac{\partial}{\partial \bar{\theta}_\alpha} - \frac{\partial}{\partial \theta_\alpha} (\not{\theta})_\alpha + \bar{\theta}_\alpha \theta_\alpha \partial^2.$$

Since the σ_μ are traceless, using the identity (A1.2) one verifies that $\bar{D}D$ can also be written as

$$\bar{D}_\alpha D_\alpha = \frac{\partial}{\partial \theta_\alpha} \frac{\partial}{\partial \bar{\theta}_\alpha} - 2(\bar{\theta}\not{\theta})_\alpha \frac{\partial}{\partial \bar{\theta}_\alpha} + \bar{\theta}_\alpha \theta_\alpha \partial^2,$$

and, therefore, in component form

$$\bar{D}_\alpha D_\alpha \Phi = 2F - 2\bar{\theta}\not{\theta}\psi + \bar{\theta}\theta \partial^2 \varphi.$$

and

$$(\bar{D}_\alpha D_\alpha)^2 = 4\partial^2.$$

Supersymmetry generators and WT identities. Supersymmetry is generated by the operators

$$Q_\alpha = \frac{\partial}{\partial \bar{\theta}_\alpha} + (\not{\theta})_\alpha, \quad \bar{Q}_\alpha = \frac{\partial}{\partial \theta_\alpha} + (\bar{\theta} \not{\theta})_\alpha,$$

which anticommute with D_α (and thus $\bar{D}_\alpha$). Then,

$$\{\bar{Q}_\alpha, Q_\beta\} = 2[\not{\theta}]_{\alpha\beta}.$$

Supersymmetry implies WT identities for correlation functions. The n -point function $W^{(n)}(p_k, \theta_k)$ of Fourier components satisfies

$$Q_\alpha W^{(n)} \equiv \left[\sum_k \frac{\partial}{\partial \bar{\theta}_\alpha^k} - i(\not{p}_k \theta^k)_\alpha \right] W^{(n)}(p, \theta) = 0.$$

To solve this equation, we set

$$W^{(n)}(p, \theta) = F^{(n)}(p, \theta) \exp \left[-\frac{i}{2n} \sum_{jk} \bar{\theta}_j (\not{p}_j - \not{p}_k) \theta_k \right], \quad (A1.5)$$

where $F^{(n)}$ is a symmetric function in the exchange $\{p_i, \theta_i\} \leftrightarrow \{p_j, \theta_j\}$. It then satisfies

$$\sum_k \frac{\partial}{\partial \bar{\theta}_\alpha^k} F^{(n)}(p, \theta) = 0,$$

that is, is translation invariant in θ space.

In the case of the two-point function, this leads to the general form

$$\begin{aligned} W^{(2)}(p, \theta', \theta) &= A(p^2) [1 + C(p^2) \delta^2(\theta' - \theta)] e^{i\bar{\theta} \not{p} \theta'} \\ &= A(p^2) [1 + C(p^2) (\bar{\theta}' - \bar{\theta})(\theta' - \theta) + i\bar{\theta} \not{p} \theta' - \frac{1}{4} p^2 \bar{\theta} \theta \bar{\theta}' \theta'] . \end{aligned} \quad (A1.6)$$

Since the vertex functions $\Gamma^{(n)}$ satisfy the same WT identities, they take the same general form.

References

- [1] L.Girardelo, M.T. Grisaru, P. Salomonson *Nucl. Phys.* B178 (1981) 331 .
- [2] A.Das and M.Kaku *Phys. Rev* D18 (1978) 4540; A. Das *Physics A* 158 (1989) 1, and references therein
- [3] A. Riotto and G. Senjanović *Phys. Rev. Lett.* 79 (1997) 349 [arXiv:hep-ph/9702319] and references therein
- [4] A. Fotopoulos and T.R. Taylor *Phys. Rev.* D59 (1999) 061701 and references therein
- [5] J.R. Espinosa *Nucl. Phys.* B475 (1996) 273 ; M. Laine *Nucl. Phys.* B481 (1996) 43 , Erratum *ibid.* B548 (1999) 637
- [6] W.A. Bardeen, M. Bander and M. Moshe, *Phys. Rev. Lett.* 52 (1983) 1188.
- [7] F.David, D.A. Kessler and H. Neuberger, *Phys. Rev. Lett.* 53 (1984) 2071 and *Nucl. Phys.* B257 (1985) 695; D. A. Kessler and H. Neuberger, *Phys. Lett.* 157B (1985) 416.
- [8] R. Gudmundsdottir, G. Rydneil and P. Salomonson, *Phys. Rev. Lett.* 53 (1984) 2529; Y. Matsubara, T. Suzuki and I. Yotsuyanagi, *Z. Phys.* C27 (1985) 599.
- [9] W.A. Bardeen, K. Higashijima and M. Moshe, *Nucl. Phys.* B250 (1985) 437, T. Reisin M.Sc. Thesis -Technion (March 1986)